

Self Organization of Edge and Internal Pedestals in a Sandpile

S. C. Chapman^{1*},

¹*Physics Dept. Univ. of Warwick, Coventry CV4 7AL, UK*

R. O. Dendy²

²*EURATOM/UKAEA Fusion Association, Culham Science Centre, Abingdon, Oxfordshire OX14 3DB, United Kingdom*

B. Hnat¹

(February 2, 2008)

Abstract

The temperature profiles of magnetically confined plasmas can display distinctive longlived pedestals at the edge and internally. Here we show that such structures can arise naturally through avalanching transport in a sandpile model. A single control parameter that is constant across the sandpile determines the occurrence and regularity of these effects, as well as the entrainment of global confinement to edge pedestal dynamics. The system dynamics indicate that the pedestals are a consequence of an inverse cascade in real space, and that self organization is necessary for their occurrence.

52.55.Dy, 52.55.Fa, 45.70.Ht, 52.35.Ra

Typeset using REVTeX

*sandr@astro.warwick.ac.uk

A distinctive feature of magnetically confined plasmas is that they can sustain local regions having very steep (indeed, almost discontinuous) temperature gradients. Pedestals in the edge temperature are a key feature of the good confinement regimes of tokamaks (“H-modes”); for a recent review see Ref. [1]. Additionally, advanced operating regimes for tokamaks have now been accessed, which involve the creation of “internal transport barriers” (ITBs) – steplike features in the temperature profiles internal to the plasma, see for example Refs. [2]- [5]. The occurrence of such structures in externally heated plasma systems that are diffuse, high temperature, and turbulent is surprising, and is a striking instance of their capacity for macroscopic self organization. Here we shall identify how similar structures arise spontaneously in the sandpile model of Ref. [6]. Because of the simplicity of this model, it is possible to formally characterize and explain the mechanisms underlying pedestal formation, and to identify links to tokamak plasma behavior. This approach is complementary to studies that employ largescale numerical simulation of tokamak plasmas, which have had some success in illuminating the subtle interplay between turbulence and bulk flows that may give rise to ITBs [5,7].

The essential ingredients of the sandpile algorithm of Ref. [6] are: (i) randomized critical gradients which must be exceeded locally before any transport can take place; (ii) central fuelling, so that sand can only leave the system by means of systemwide avalanches; and (iii) its only distinctive feature, namely a characteristic lengthscale L_f for fast redistribution which may be considered a proxy for turbulent transport, as discussed in Ref. [8]. Evidence for avalanche-type transport from tokamak experiments [9,10] and numerical simulations [11,12] provides growing support for the applicability in some circumstances of the sandpile paradigm introduced into fusion plasma physics in recent years [13]- [17]. Particularly relevant to the present study are observations and analysis [18,19] of edge plasma turbulence in a range of magnetically confined plasmas. These suggest that edge plasma turbulence – the environment in which H-mode edge pedestals form, which then coexist with the turbulence – self organizes into a critical state, independent of the size and plasma characteristics of the devices considered. It is known [8] that, depending on the value of the control parameter L_f , the statistical behavior of the sandpile model of Ref. [6] displays features reminiscent of enhanced confinement phenomenology in tokamaks. These include the time averaged height profiles, which possess edge pedestals in the good confinement regime; furthermore the frequency of systemwide avalanches resulting in mass loss scales with stored sandpile energy in the same way as the frequency of edge localized modes (ELMs) scales with stored energy in tokamaks, see Fig.6 of Ref. [8]. We emphasize that L_f is kept constant across the sandpile in any given computational run, so that the critical gradient and redistribution rules are identical at every cell of our sandpile, apart from the initial and final cells. It follows that where pedestals arise, they represent a true emergent phenomenon. In this respect our approach differs essentially from that of Refs. [13,17], where a local transport barrier is specified at the outset, by declaring the critical gradient and redistribution rules to be different for a specific local group of cells.

The three dimensional plot of Fig.1 shows sandpile height as a function of position as time evolves, for the good confinement regime [8] with $L_f = 50$ in a 512 cell system. It displays two distinct phases. First, there is a relaxation phase where the sandpile profile is smooth down to the self organized edge pedestal, except within a distance L_f of the core where fueling has a continual local effect. During the relaxation phase mass loss occurs via many

systemwide avalanches closely spaced in time, which carry sand over the sandpile boundary. The relaxation phase terminates with a final systemwide avalanche, after which the growth phase begins. The growth phase is characterized by a stationary edge pedestal which resides at the outermost cell of the sandpile. As time progresses, additional pedestals (localized regions of steep gradient just below critical) form successively at positions increasingly close to the core of the sandpile, with average separation $\sim L_f$. Each of these is generated at positions where (outward propagating) major internal avalanches have come to rest. The location of the most recently formed (and therefore innermost) internal pedestal propagates inward during the growth phase.

Figure 2 is motivated by simultaneous multichannel measurements of tokamak temperature profiles in the presence of ITBs; see for example Fig.3 of Ref. [3] and Fig.1 of Ref. [5], and also the results of numerical simulations shown in Fig.2 of Ref. [5]. Figure 2 shows sandpile height at different positions from the edge to the centre. Just over two growth and relaxation phases are shown. The successive formation of internal pedestals is reflected in a stepwise increase in height at any given point during the growth phases. Using this diagnostic, evidence for stepwise increments is clearest in the region of the sandpile that lies between the core and the edge. Since the internal pedestals form at locations increasingly close to the core as the growth phase proceeds, the points within the sandpile that are most affected by the formation of these pedestals are those that are closer to the core of the sandpile. Points nearer the edge are only affected by the formation of the first few internal pedestals during the early growth phase. The results shown in Fig.2, which bear some resemblance to the experimental and numerical results of Refs. [3,5], emerge naturally from the dynamics of the sandpile during the growth phase of its good confinement regime. Central to this structure is the unexpected capacity of this sandpile to organize persistent steep pedestals both at the edge and internally.

The role of these internal pedestals and their relationship to the edge pedestal is highlighted in Fig.3: here all cells at which the gradient exceeds $z_c/2$ are marked by black points, while all other cells are left blank. Figure 3(a) ($L_f = 50$) follows five of the growth and relaxation cycles shown in Fig.2. The edge pedestal is visible close to the sandpile boundary in both the relaxation and the growth phases. Its time behaviour is essentially regular and, as we shall see, orders the structure internal to the sandpile. For any $L_f < N/4$, the location of each internal pedestal is fixed during a given growth phase, so that they persist as distinguishable features of the time averaged phenomenology of the sandpile. In contrast, Fig.3(b) shows the behavior for the poor confinement regime with $L_f = 250 > N/4$. This regime corresponds to a self organized critical (SOC) state, see Ref. [8] and below. Although the sandpile successively fills and empties, it does so in an irregular manner. Pedestals can be seen both at the edge and internal to the sandpile, but these are no longer organized in a coherent pattern.

A resilient edge pedestal arises for all L_f ; the pedestal is steep, indeed unresolved, in that the entire change in height occurs between neighboring cells. The location of the edge pedestal is strongly time dependent for large L_f , whereas for small L_f it is confined to a region close to the outermost cell of the sandpile. The time averaged profile in the edge region therefore depends strongly on L_f , and the steadiest edge pedestal corresponds to small L_f and good confinement. Following each avalanche, in our algorithm the value of the critical gradient z_c is randomised about a mean value at all cells that participated in

the avalanche. Nevertheless the gradient at the edge pedestal remains always close to, but just below, the critical value z_c as shown in Fig.4. Elsewhere in the sandpile (for example at its midpoint, see Fig.4), except where internal pedestals arise, the gradient is well below z_c . The internal pedestals appear as barriers to transport: despite their apparent fragility against avalanching (gradient z close to z_c), no sand passes through either the edge or the internal pedestals until the final avalanche that terminates the growth phase.

The physical mechanisms and principles underlying the self organization of the edge pedestal and multiple internal pedestals that arise in the sandpile model of Refs. [6,8] are therefore of considerable interest. As a first step, we confirm the hypothesis of Ref. [8] that the good confinement regime (small L_f) corresponds to low dimensional behavior. In Fig.5 the position of the last occupied cell at time t is plotted against that at time $t + \tau$, where $\tau = 50$, for runs with $L_f = (a)50$, $(b)150$ and $(c)250$ in a 512 cell system. This is an example of phase space reconstruction, achieved here by embedding [20]. Figure 5(a) shows low dimensional system dynamics that repeatedly follow a simple limit cycle (attractor) around a restricted region of the reconstructed phase space. This implies that the large number of cells in the sandpile have self organized: their collective dynamics are encapsulated by a small number of dynamical variables. Once L_f is increased to 150 (Fig.5(b)), the simple limit cycle seen in Fig.5(a) bifurcates, and more stochastic behavior is seen in Fig.5(c) ($L_f = 250$). Thus increasing complexity of the phase space portrait correlates with deterioration of confinement [8]. The lowest confinement regime corresponds to selfsimilar avalanche statistics. This is associated with a nontrivial fixed point in the space of the parameter used [21] to perform rescaling under the renormalization group procedure, corresponding to behavior that is both self organized and critical [22]. Importantly, global relaxation of the sandpile is ultimately achieved by large (systemwide) avalanches for all values of L_f . When L_f is of order the system size N , systemwide avalanches are straightforwardly propagated: because $L_f \sim N$, no characteristic scale is imposed by the redistribution process and the dynamics are selfsimilar and in SOC [21]. Conversely, when L_f is significantly distinct from the system size (found empirically to be $L_f < N/4$), there is scale breaking. This leads to broken power law avalanche statistics [6], and the system is no longer in SOC. By separating the characteristic lengthscales we also effectively separate the longterm growth-relaxation timescale from the time interval between systemwide avalanches. In the SOC regime, such a distinction is not possible. Thus, in the good confinement regime (when $L_f < N/4$) the requirement for self organization is satisfied. The feature evolving on the slow timescale, namely the position of the edge pedestal, is sufficient to determine the details of the internal dynamics. It organizes the sequence of events leading to the successive formation of internal edge pedestals and the time variation of total energy (sand) in the system: in short, the sandpile is entrained to its edge. The characteristic signature of the onset of self organization is low dimensional dynamics, seen in Figs.4($L_f = 50$ traces) and 5(a), as opposed to the irregular time evolution and selfsimilar statistical properties of the system when self organised and critical (Figs.4($L_f = 250$ traces) and 5(c)).

The confinement physics of our sandpile model offers a robust framework in which a distinctive structure of edge and internal pedestals (previously known only from tokamak plasmas) arises naturally. There is only one control parameter, L_f/N , which can be considered as a proxy for the lengthscale of turbulent transport, normalized to system size. Provided that this lengthscale is sufficiently short, the underlying inverse cascade in real

space gives rise to persistent, marginally subcritical profile steps whose formation point propagates inwards in the growth phase. Furthermore the self organized edge pedestal is continuously present in all phases of the sandpile evolution, and positions itself exactly at the sandpile boundary throughout the growth phase. These results are sufficient to indicate that some of the distinctive edge and internal pedestal phenomenology seen in tokamak plasmas can also arise in a simpler idealized confinement system, and that they may be linked to the observed avalanching transport phenomena. This suggests that quantitative studies to elucidate the balance between diffusive transport and avalanching transport in tokamaks would be fruitful. The results further suggest a test of the depth of the physical analogy that we have found, as follows. If the analogy is deep, there will exist one or a few dimensionless control parameters, linked to the properties of the turbulent transport, that entirely determine the key features of the confinement phenomenology – regular or irregular – of tokamak plasmas that we have cited here. For example, these parameters would control the extent to which global confinement is entrained to edge pedestal dynamics. The search for such parameters, for example by further application of the techniques of nonlinear time series analysis to edge plasma measurements as initiated in Refs. [18,19], is potentially highly rewarding.

ACKNOWLEDGMENTS

We are grateful to George Rowlands and Jack Connor for comments and suggestions. SCC was supported by a PPARC lecturer fellowship, ROD by Euratom and the UK DTI, and BH by HEFCE.

REFERENCES

- [1] J. Hugill, Plasma Phys. Control. Fusion **42**, R75 (2000).
- [2] E. J. Synakowski, Plasma Phys. Control. Fusion **40**, 581 (1998).
- [3] K. H. Burrell *et al.*, Plasma Phys. Control. Fusion **40**, 1585 (1998).
- [4] G. D. Conway *et al.*, Phys. Rev. Lett. **84**, 1463 (2000).
- [5] J. E. Kinsey, G. M. Staebler, K. H. Burrell, M. E. Austin, and R. E. Waltz, Phys. Rev. Lett. **86**, 814 (2001).
- [6] S.C. Chapman, Phys. Rev. E **62**, 1905 (2000).
- [7] B. N. Rogers, W. Dorland, and M. Kotschenreuther, Phys. Rev. Lett. **85**, 5336 (2000).
- [8] S. C. Chapman, R. O. Dendy, and B. Hnat, Phys. Rev. Lett. **86**, 2814 (2001).
- [9] T. L. Rhodes *et al.*, Phys. Lett. A **253**, 181 (1999).
- [10] P. A. Politzer, Phys. Rev. Lett. **84**, 1192 (2000).
- [11] X. Garbet and R. Waltz, Phys. Plasmas **5**, 2836 (1998).
- [12] Y. Sarazin and P. Ghendrih, Phys. Plasmas **5**, 4214 (1998).
- [13] D. E. Newman, B. A. Carreras, P. H. Diamond, and T. S. Hahm, Phys. Plasmas **3**, 1858 (1996).
- [14] B. A. Carreras, D. Newman, V. E. Lynch, and P. H. Diamond, Phys. Plasmas **3**, 2903 (1996).
- [15] R. O. Dendy and P. Helander, Plasma Phys. Control. Fusion **39**, 1947 (1997).
- [16] S. C. Chapman, R. O. Dendy, and G. Rowlands, Phys. Plasmas **6**, 4169 (1999).
- [17] H. R. Hicks and B. A. Carreras, Phys. Plasmas **8**, 3277 (2001).
- [18] B. A. Carreras *et al.*, Phys. Rev. Lett. **80**, 4438 (1998).
- [19] M. A. Pedrosa *et al.*, Phys. Rev. Lett. **82**, 3621 (1999).
- [20] E. Ott, *Chaos in Dynamical Systems* (Cambridge Univ. Press, 1993), p 93.
- [21] S. W. Y. Tam, T. S. Chang, S. C. Chapman, and N. W. Watkins, Geophys. Res. Lett. **27**, 1367 (2000).
- [22] J. P. Sethna, K. A. Dahmen, and C. R. Myers, Nature, **410**, 242 (2001).

Captions

FIG 1: Three dimensional view of the sandpile height profile for $L_f = 50$. Time evolves along the partially hidden axis. One relaxation phase and one subsequent growth phase are shown.

FIG 2: Local height of sand as a function of time at different locations in the sandpile relative to the centre cell $n = 1$: (a) $n=20$, (b) $n=65$, (c) $n=100$, (d) $n=125$, (e) $n=150$ and (f) $n=300$. System size $N = 512$, control parameter $L_f = 50$.

FIG 3: Location of cells where the value of the local gradient exceeds $z_c/2$ for (a) $L_f = 50$ and (b) $L_f = 250$. Zero corresponds to the apex where fueling occurs. The edge pedestal is visible as the uppermost trace.

FIG 4: Local gradient normalized to the local value of z_c for cells at the edge (Δ_e , upper traces, value close to unity) and halfway into the sandpile (Δ_{mid} , lower traces, value close to zero), for $L_f =$ (top) 250, (centre) 150, (bottom) 50.

FIG 5: Phase space reconstruction of the dynamics of the edge position $ep(t)$. Plotted are coordinates $ep(t)$ versus $ep(t + \tau)$ for $\tau = 50$ and $L_f =$ (a) 50, (b) 150 and (c) 250. The system dynamics explore a larger region of the phase space with increasing values of L_f . The topology shown is insensitive to the value of τ in the range of interest.















